Structure

Graphical Abstract



Authors

R. Elliot Murphy, Alexandra B. Samal, Jiri Vlach, Jamil S. Saad

Article

Correspondence saad@uab.edu

saad@uab.edu

In Brief

Murphy et al. devised new approaches to prepare the HIV-1 gp41CT protein, solved the structure by NMR methods, characterized its interactions with the membrane, and provided a preferred topology of the protein when bound to the membrane.

Highlights

- Devised new approaches to generate HIV-1 gp41CT protein via recombinant techniques
- Determined the NMR structure of gp41CT in micellar solution
- Characterized gp41CT interaction with the membrane
- Provided a preferred topology of gp41CT bound to the membrane



Structure Article

Solution Structure and Membrane Interaction of the Cytoplasmic Tail of HIV-1 gp41 Protein

R. Elliot Murphy,^{1,2} Alexandra B. Samal,^{1,2} Jiri Vlach,¹ and Jamil S. Saad^{1,3,*}

¹Department of Microbiology, University of Alabama at Birmingham, Birmingham, AL 35294, USA

²These authors contributed equally

³Lead Contact

*Correspondence: saad@uab.edu

https://doi.org/10.1016/j.str.2017.09.010

SUMMARY

The cytoplasmic tail of gp41 (gp41CT) remains the last HIV-1 domain with an unknown structure. It plays important roles in HIV-1 replication such as mediating envelope (Env) intracellular trafficking and incorporation into assembling virions, mechanisms of which are poorly understood. Here, we present the solution structure of gp41CT in a micellar environment and characterize its interaction with the membrane. We show that the N-terminal 45 residues are unstructured and not associated with the membrane. However, the C-terminal 105 residues form three membrane-bound amphipathic α helices with distinctive structural features such as variable degree of membrane penetration, hydrophobic and basic surfaces, clusters of aromatic residues, and a network of cation- π interactions. This work fills a major gap by providing the structure of the last segment of HIV-1 Env, which will provide insights into the mechanisms of Gag-mediated Env incorporation as well as the overall Env mobility and conformation on the virion surface.

INTRODUCTION

The envelope glycoprotein (Env) of human immunodeficiency virus type 1 (HIV-1) is synthesized as a 160-kDa precursor in the rough endoplasmic reticulum, where it is glycosylated, and subsequently cleaved in the Golgi apparatus to the form of the surface (gp120) and transmembrane (gp41) subunits (reviewed in Checkley et al., 2011). The gp41 subunit comprises a fusogenic ectodomain, a transmembrane domain, and a C-terminal cytoplasmic tail (gp41CT). For the past two decades, research has mainly focused on elucidating the mechanisms of the gp120/ gp41 trimer binding to host receptor, membrane fusion, and evasion of recognition by the immune system. It is established that binding of gp120 to the CD4 receptor and CCR5/CXCR4 co-receptors triggers large conformational changes in the gp41 ectodomain, allowing the viral membrane to fuse with the host membrane (reviewed in Merk and Subramaniam, 2013). The membrane-proximal external region (MPER) of the HIV-1 gp41 protein, which forms an α -helical trimer (Buzon et al., 2010), precedes the transmembrane domain and is a key target for broadly neutralizing monoclonal antibodies elicited during human infection (Alam et al., 2009). Current models suggest that receptor binding leads to the exposure of the gp41 fusion peptide, which anchors to the target cell membrane to produce an intermediate, pre-hairpin state bridging the two membranes.

One of the least understood phenomena in retroviral assembly is the mechanism by which the Env protein is recruited and incorporated into virus particles. There is mounting evidence that gp41CT plays a functional role in Env incorporation in physiologically relevant cell types (Akari et al., 2000; Murakami and Freed, 2000b). Deletion of gp41CT strongly reduces Env incorporation in permissive cell lines but has only minor effect in non-permissive cell lines (Freed and Martin, 1995, 1996; Murakami and Freed, 2000b). The CT domain is remarkably long (150 amino acids) for most lentiviruses but significantly shorter (~20–40 amino acids) for other retroviruses (Checkley et al., 2011). The biological implication of the variable length and its effect on Env incorporation are not well understood.

The assembly of HIV-1 particles is an orchestrated process that is driven by transport of Gag polyproteins to the plasma membrane (PM) (reviewed in Freed, 2015). Gag-membrane binding is mediated by the myristoylated N-terminal matrix (MA) domain. HIV-1 Gag targeting to the PM is dependent on phosphatidylinositol 4,5-bisphosphate (PI(4,5)P2) (Chukkapalli and Ono, 2011; Ono et al., 2004), which binds directly to the MA domain (Anraku et al., 2010; Mercredi et al., 2016; Saad et al., 2006; Shkriabai et al., 2006; Vlach and Saad, 2013). Based on genetic, in vivo, and biochemical evidence, early studies suggested that Env incorporation is mediated by interactions between the MA domain of Gag and gp41CT (Cosson, 1996; Dorfman et al., 1994; Freed and Martin, 1995, 1996; Yu et al., 1992). Freed and co-workers have recently provided biochemical evidence that MA trimerization is an obligatory step in the assembly of infectious HIV-1 virions and demonstrated a correlation between loss of MA trimerization and loss of Env incorporation (Tedbury et al., 2016). It has also been shown that Gag assembly induced the aggregation of small Env clusters into larger domains that were completely immobile (Roy et al., 2013). Truncation of the CT domain abrogated Gag's ability to induce Env clustering and restored Env mobility at assembly sites (Roy et al., 2013). Super-resolution microscopy data indicated that recruitment of HIV-1 Env to viral assembly sites is dependent on gp41CT (Muranyi et al., 2013). Spearman and co-workers also demonstrated that Rab11-family interacting

Α

707 RVRQGYSPLS FQTHLPIPRG PDRPEGIEEE GGERDRDRSI 747 RLVNGSLALI WDDLRSLCLF SYHRLRDLLL IVTRIVELLG 787 RRGWEALKYW WNLLQYWSQE LKNSAVNLLN ATAIAVAEGT 827 DRVIEVLQAA YRAIRHIPRR IRQGLERILL



Figure 1. NMR Spectra of the Two Independent gp41CT Domains (A) Amino acid sequence of the HIV-1 gp41CT protein (pNL4-3 isolate, HXB2 numbering). Sequences of gp41CT_N and gp41CT_C are highlighted in blue and black, respectively.

(B) 2D ¹H-¹⁵N HSQC spectrum obtained for gp41CT_N (200 μ M) at 32°C. (C) 2D ¹H-¹⁵N HSQC spectrum obtained for gp41CT_C (300 μ M) at 50°C. Tryptophan side-chain signals are colored in magenta (inset). Dashed lines indicate side-chain amide signals.

protein 1C (FIP1C) is required for CT-dependent incorporation of Env into HIV-1 particles (Qi et al., 2013, 2015), and identified a tyrosine-based motif (Y795/W796) as critical in mediating celltype-dependent Env incorporation (Qi et al., 2015). FIPs are effectors of Rab11 GTPases that mediate sorting of cargo from the endosomal recycling compartment to the PM (Hales et al., 2001). Altogether, these lines of evidence strongly suggest a role of gp41CT in Gag-Env co-localization and incorporation of Env into virus particles.

Despite the evidence for a Gag-mediated Env incorporation process, structural or biophysical evidence for direct gp41CT-MA interaction is still unavailable. A major barrier to characterizing a potential gp41CT-MA interaction by structural tools has been the unavailability of a recombinant gp41CT protein. A structural and functional topology of the gp41CT domain has been proposed based on sequence analysis and biophysical characterization of short peptide fragments (Boscia et al., 2013; Costin et al., 2007; Steckbeck et al., 2010, 2013). A model of gp41CT has been proposed in which a portion of the protein appears to be associated with the membrane (Costin et al., 2007; Steckbeck et al., 2010). The membraneassociated domain contains three lentivirus lytic peptide motifs called LLP2, LLP3, and LLP1, which are highly conserved not only among HIV-1 strains but also among HIV-2, simian immunodeficiency virus (SIV), and equine infectious anemia virus (EIAV) (Srinivas et al., 1992; Steckbeck et al., 2011). However, an exact protein topology is still controversial. Conventional models based on primary sequence analysis and biochemical assays have placed the entirety of CT inside the cytoplasm of the cell/interior of the virus (Steckbeck et al., 2010). Alternative models based on the presence of an immunogenic phenotype (Kennedy epitope, KE) have been proposed, which postulate the existence of a transmembrane region within gp41CT allowing for the extracellular exposure of the immunogenic region (Cleveland et al., 2003; Hollier and Dimmock, 2005; Steckbeck et al., 2010). Structural characterization of the gp41CT protein is therefore critical for determining its precise topology and the functional role in Gagmediated Env incorporation.

Here, we devised new approaches that allowed for successful preparation of the gp41CT protein in micellar solution and determination of the structure by nuclear magnetic resonance (NMR) methods. We show that the N-terminal region (45 amino acids) of gp41CT lacks a regular secondary structure and is not associated with membrane. The C-terminal domain (105 amino acids), however, forms three consecutive amphipathic α helices that are tightly associated to membrane. This study solves a long-standing challenge and provides new insights into the mechanisms of Gag-mediated Env incorporation as well as the overall Env mobility and conformation on the virion surface.

RESULTS

Two Independent Domains of gp41CT

We have devised a strategy to produce the gp41CT protein via recombinant techniques, which involved expression, purification, and screening for the proper membrane mimetics to solubilize the protein and allow for detection of NMR signals. Details on gp41CT preparation and biochemical and biophysical properties are described in STAR Methods. The gp41CT protein was, however, prone to proteolysis during expression and purification resulting in a soluble peptide comprising residues 707–751 (gp41CT_N) and an insoluble fragment comprising residues 752–856 (gp41CT_C) (Figure 1A). To facilitate their biochemical, biophysical, and structural characterization, we have expressed and purified the two fragments independently. As described below, we show that the two domains adopt structures that are very similar to those within the intact protein.

Characterization of gp41CT_N

The gp41CT_N fragment was soluble at concentrations >1 mM in the absence of detergents, which enabled full structural and biophysical characterization. On a gel filtration column, gp41CT_N elutes as a single peak at 17.9 mL (Figure S1), indicating a homogeneous sample. When compared with known protein markers, the molecular mass of the $gp41CT_N$ fragment corresponds to a \sim 13 kDa species (Figure S1). The 2D ¹H-¹⁵N heteronuclear single quantum coherence (HSQC) spectrum of ¹⁵N-labeled gp41CT_N shows a narrow dispersion of the amide proton resonances, suggesting a lack of ordered structure (Figure 1B). The far-UV circular dichroism (CD) spectrum of gp41CT_N displays a negative band at ${\sim}200$ nm, consistent with a random coil conformation (Figure S2). No changes are observed in the CD spectrum of $gp41CT_N$ in the presence of n-dodecylphosphocholine (DPC), demonstrating that protein does not associate with membrane. Proton, carbon, and nitrogen chemical shifts of gp41CT_N were used to predict order parameters and secondary structure content in TALOS+ (Shen et al., 2009), confirming that gp41CT_N lacks any regular secondary structure (Figure S2). Nuclear Overhauser effect spectroscopy (NOESY) data also indicated that the protein lacks a regular secondary structure. No intermolecular NOEs indicative of a protein oligomer have been detected, confirming that gp41CT_N is a monomer and that the smaller than expected elution volume obtained from gel filtration data is likely due to the shape and disordered conformation of the molecule. Characterization of gp41CT_c

Preparation and membrane reconstitution of the gp41CT_C fragment was conducted in a manner that is essentially identical to that of the full-length gp41CT protein (STAR Methods). The best results regarding protein reconstitution, stability, and quality of NMR data were achieved in DPC micellar solution. A gel filtration assay of gp41CT_C incorporated in DPC micelle shows a single peak at 15.2 mL, which corresponds to \sim 65 kDa species (Figure S1). Similar to the gp41CT protein, the apparent molecular mass of gp41CT_C in DPC micelles as determined by gel filtration is significantly larger. gp41CT_C in DPC produced highquality NMR spectra at 50°C, which enabled a full structural characterization as described below. CD spectra obtained for gp41CT_C in DPC micelles at 35 and 50°C (Figure S2) are virtually identical, demonstrating that the protein structure was not compromised at 50°C.

We also examined whether the gp41CT_C protein can produce high-quality NMR data in dimyristoylphosphatidylcholine (DMPC)/dihexanoylphosphatidylcholine (DHPC) bicelles and whether the NMR spectra are similar to those obtained in DPC micelles. Although the 2D ¹H-¹⁵N signals in the HSQC spectrum of gp41CT_C in DMPC/DHPC bicelles were relatively broad compared with those observed in DPC micelles, their positions were remarkably similar (Figure S3), demonstrating that gp41CT_C maintains a similar structure in both membrane environments. A question may arise whether the structural properties of the isolated gp41CT_N and gp41CT_C proteins are similar to the corresponding domains within the full-length gp41CT protein. To answer this question, we compared the ¹H-¹⁵N HSQC spectrum of gp41CT with those of the isolated gp41CT_N and gp41CT_C domains under identical buffer conditions in DPC micelles. As shown in Figure S3, the ¹H-¹⁵N resonances of gp41CT_N and gp41CT_C were at very similar positions to the corresponding signals in the spectrum obtained for the full-length gp41CT protein, which demonstrates that the two domains adopt structures that are very similar to those in the intact protein. Importantly, NMR data analysis of gp41CT_C was facilitated by the absence of strong and overlapping gp41CT_N signals allowing for full structural characterization of the protein.

NMR Signal Assignment of gp41CT_c

The majority of backbone amide NMR signals were assigned with the exception of L771, R772, L776, I843, R845, R846, and 1847 (Figure 1C). Overall, more than 80% of $gp41CT_{C}$ ¹H, ¹³C, and ¹⁵N resonances were assigned, with the exception of L771, R845, and R846 side chains, which could not be resolved due to severe signal overlaps or signal broadening. For P844, only very broad signals of C^{δ} and C^{γ} groups were detected, suggesting that the lack of signals within this region (P844-R846) was due to an intermediate conformational exchange. The chemical shift assignments for gp41CT_C were used to predict its order parameters and secondary structure content in TALOS+ (Figure S2). The TALOS + data indicated that gp41CT_C contains three α -helical motifs spanning residues 753–785, 790–823, and 827-853. These motifs were named LLP2, LLP3, and LLP1, respectively (Figure 2) (Steckbeck et al., 2010). In the two short linkers, the α -helical character appeared to be either reduced (residues 786-789 between LLP2 and LLP3) or not present (residues 824-826 between LLP3 and LLP1). Taken together, our data indicated that $gp41CT_N$ lacks an ordered structure and is not associated with membrane. gp41CT_C, however, is tightly associated with membrane and formed three consecutive α helices connected by short linkers. The structures and dynamics of the two domains appeared to be completely independent of each other.

Structure Determination of gp41CT_c

Sequential and medium-range NOEs obtained from 3D ¹⁵N-edited NOESY-HSQC and ¹³C-edited heteronuclear multiple-quantum correlation (HMQC)-NOESY spectra contained an extensive network of characteristic a-helical contacts. No long-range intraor intermolecular NOEs indicative of a tertiary structural fold or formation of oligomer of the gp41CT_C protein were detected. A total of 492 NOEs were used to calculate the structure (Table 1). The calculated ensemble of 20 lowest-energy structures showed a good convergence and correspondingly low positional rootmean-square deviation (RMSD) values within the structured regions (Figures 3A and S3 and Table 1). The gp41CT_C protein adopted an α -helical conformation with clearly defined α helices spanning residues 753-785, 790-823, 827-841, and 847-853 (Figure 3). The linker regions (residues 786–789 and 824–826) were mostly unrestrained during structure calculations and the three a helices in the resulting structures adopted pseudorandom orientations with respect to each other, limited only by steric factors (Figure S3). Therefore, the bending observed between the helical motifs is variable in the 20 calculated structures. Because signals corresponding to residues P844, R845, and R846 were undetectable, this region was also not restrained during calculation resulting in a small break in the LLP1 helical structure. Of note, previous NMR studies conducted in the presence of 2,2,2-trifluoroethanol on an LLP1 peptide spanning residues 827-853 have shown that the region involving residues 842–845 is highly flexible and adopts a type-II β -turn at P844



Figure 2. Secondary Structure and Helical Wheel Diagrams of the gp41CT_C Protein (A) Secondary structure representation of the gp41CT protein based on the NMR data.

(B) Helical wheel diagrams of the LLP2, LLP3, and LLP1 motifs. Amino acid sequences are plotted clockwise. Hydrophobic and aromatic residues are represented by light and dark green squares, respectively. Polar residues are shown as yellow circles, while positively and negatively charged residues as shown as blue and red pentagons, respectively. The helical wheels are oriented so that their hydrophobic moments (indicated in the centers) point upwards. Helical wheels were generated via a modified script obtained from http://rzlab.ucr.edu/scripts/wheel/.

and R845 (Sham et al., 2008; Yuan et al., 1995). While we could not obtain any structural data for this region, the observed broadness of P844 methylene signals is consistent with a highly flexible region. Another structural feature of gp41CT_C is the presence of strong NOE contacts between aromatic rings of tyrosine, tryptophan, or phenylalanine that are in the *i* position and sidechain groups of arginine (H^{δ}) or lysine (H^{ϵ}) in the *i* + 4 position, suggesting the presence of a network of cation- π interactions (Gallivan and Dougherty, 1999). Such NOEs were detected for F766–R770, Y768–R772, W790–K794, and Y737–R741, but not for the W757–R761 pair. Interestingly, a strong association also exists for W797–Q801 and W802–E806 pairs given the presence of strong NOEs between the respective side chains and significant upfield shifts of the Gln and Glu resonances.

A striking feature gleaned from the gp41CT_C structure was the extensive distribution of hydrophobic surface (Figures 3B and 3C). Hydrophobic and aromatic residues comprise >50% of the total gp41CT_C protein sequence. The hydrophobic surface is a characteristic feature of the amphipathic gp41CT_C and indicates that the interaction interface with the membrane is extensive (see below). An unusual feature of the gp41CT_C sequence was the clustering of six aromatic residues at the beginning of helix LLP3 (Figure 3D). Several of these aromatic residues were implicated in the functional role of gp41CT in Env incorporation and infectivity (Lambele et al., 2007; Murakami and Freed, 2000a; Qi et al., 2015; Qi et al., 2013). As shown in Figure 2B, these residues form three pairs with differential localization in the amphipathic structure. Residues W790 and W797 are located on the hydrophobic-hydrophilic interface and, as described above, are engaged in the cation- π and anion- π interactions, whereas W796 and W803 are located on the hydrophobic side of the α helix. On the other hand, Y795 and Y802 are both localized on the polar side. Interestingly, Y795/W796 has been shown to be a critical motif mediating cell-type-dependent Env incorporation (Qi et al., 2015).

Another characteristic feature of the gp41CT_C protein was the unusually high concentration of arginine residues in LLP2 and LLP1 (Figures 3C and 3E). Recent studies have shown that while conservative substitution of LLP2 Arg residues displayed wild-type phenotypes, similar substitution of LLP1 Arg residues resulted in a significant impairment of Env expression, fusogenicity, and incorporation, as well as virus replication (Kuhlmann et al., 2014). As shown in Figures 2B and 3E, six arginine residues are concentrated on the polar side of LLP1 which generates a basic surface in the C terminus of gp41CT_C (Figures 3C and 3E).

Taken together, our structural data revealed that gp41CT_C adopted an α -helical conformation with unique characteristics such as the extensive hydrophobic surface, clustering of aromatic residues in LLP3, unusually high number of cation- π interactions, and formation of a basic patch on the polar side of LLP1. Of note, our structure is among the largest membrane-associated systems studied by NMR spectroscopy, second only to the reported structure and dynamics of micelle-bound α -synuclein, a 140-amino acid protein implicated in Parkinson's disease (Ulmer and Bax, 2005; Ulmer et al., 2005).

Protein-micelle Interactions

The structure of the micelle-bound $gp41CT_{C}$ protein revealed a large hydrophobic surface that is predisposed to interact with membrane (Figures 2B and 3B). We employed three

CellPress

Table 1. NMR Refinement Statistics of the 20 Model Ensemble of $gp41CT_{c}$

<u></u>	
NMR-Derived Restraints	
¹ H- ¹ H distance restraints	492
Intraresidue	115
Sequential ($ i - j = 1$)	156
Medium range (1 < $ i - j \le 4$)	221
Long range (4 < $ i - j $)	0
Backbone H bonds (4/H bond)	77
Distance restraints per refined residue	8.9
Torsion angles	198
Restraint Violations (Mean ± SD)	
Average maximum upper distance violations (Å)	0.20 ± 0.02
Average maximum van der Waals violations (Å)	0.19 ± 0.04
Average maximum torsion angle violations (Å)	0.87 ± 0.52
Cyana target function (Å ²)	0.65 ± 0.14
Residue Distribution in Ramachandran Plot (%)	
Most favored regions	97.0
Additional allowed regions	2.6
Generously allowed regions	0.4
Disallowed regions	0.0
Structure Convergence (Å) ^a (Mean ± SD)	
Backbone N, C ^{α} , C ^{\prime} atoms RMSD	
Residues 754–785	0.62 ± 0.24
Residues 791–820	0.26 ± 0.10
Residues 827–842	0.22 ± 0.07
Residues 848–855	0.15 ± 0.03
All heavy atoms RMSD	
Residues 754–785	1.14 ± 0.16
Residues 791–820	0.77 ± 0.10
residues 827-842	1.09 ± 0.13
Residues 848–855	1.26 ± 0.15
2	

^aStructures were superimposed with the representative model and RMSDs against an average structure calculated for the indicated residue ranges.

complementary approaches to probe the depth of membrane insertion by gp41CT_C. First, we obtained 2D NOESY, 3D ¹³C-edited HMQC-NOESY, and ¹³C-half-filtered/¹³C-edited NOESY-HSQC data and identified numerous intermolecular NOE cross-peaks between DPC and the side chains of hydrophobic and aromatic residues (Figure 4). The methylene resonances of the DPC chain exhibited strong intermolecular NOE correlations to the side chains of W757, F766, Y768, W790, W797, W803, V812, V832, Y837, and I840. Residues Y795 and Y802 exhibited only weak NOEs with the DPC micelle interior groups but relatively strong NOEs with the DPC polar head (Figure 4), confirming their exposure on the micelle surface. As expected, strong NOEs were also observed with the side chains of the bulk of Leu, Ile, and Ala residues (Figure 4). In general, NOEs between the side chains of most hydrophobic/aromatic residues and the polar

head of DPC were virtually nonexistent or very weak, indicating a significant penetration of the protein in the interior of DPC micelles.

In the second approach, we assessed the depth of membrane insertion by utilizing two paramagnetic spin-labeled "depth" probes, 5-doxyl stearic acid (5-DSA) and 16-doxyl stearic acid (16-DSA). Both molecules are incorporated into DPC micelles and cause severe broadening of the NMR signals of nuclei that are within \sim 10–12 Å of the paramagnetic center (Jarvet et al., 1997; Papavoine et al., 1994). For 16-DSA, only the intensities of resonances buried in the micellar core are dampened. Addition of 16-DSA to ¹⁵N- or ¹³C-labeled gp41CT_C samples in DPC micelles caused a substantial reduction of intensities of numerous signals in the 2D ¹H-¹⁵N and ¹H-¹³C HSQC spectra, respectively. Quantification of the line broadening of the amide resonances due to the paramagnetic relaxation enhancement clearly showed that the effect is widespread throughout the entire gp41CT_C protein (Figure S4). Likewise, significant reduction of signal intensities was observed upon addition of 5-DSA to gp41CT_C in DPC micelles (Figure S4). Overall, signal dampening caused by 16-DSA was larger for the majority of residues compared with the 5-DSA analog, indicating that the gp41CT_C protein was significantly inserted in the interior of the micelle core.

In the third approach, we probed the regions that are most exposed to solvent by titrating gadodiamide, which is excluded from the hydrophobic interior of the micelle. Paramagnetic gadodiamide induces distance-dependent broadening of NMR signals of residues that are solvent exposed or near the aqueous environment. We monitored the gadodiamide-induced line broadening in the ¹H-¹⁵N HSQC spectra obtained for gp41CT_C. Although there was a moderate widespread reduction in signal intensity at high concentration of gadodiamide, a substantial decrease in the intensity of ¹H-¹⁵N resonances was observed for residues 825-827 (Figure S4), indicating exposure of the linker between LLP3 and LLP1 to solvent. Other residues such as R788 (located in the linker between LLP2 and LLP3), E791, Y802, S810, A817, Q831, and G850 were also affected but to a lesser extent, suggesting that these regions are transiently exposed or less buried in micelles. Taken together, our NMR data have shown that gp41CT_C interacted extensively with the micelle and that numerous hydrophobic residues were buried in the interior of the membrane.

DISCUSSION

Several points have emerged from this work. Most importantly, this study fills a major gap by providing the structure of the last remaining segment of HIV-1 Env, which will likely provide insights into the mechanisms of Env incorporation as well as the overall Env mobility and conformation on the virion surface. We show that gp41CT consists of an unstructured N-terminal domain (residues 707–751) and a membrane-bound C-terminal domain (residues 752–856). A recent NMR structure of the transmembrane segment of gp41 revealed that residues 707–710 (RVRQ) located at the C terminus of the transmembrane α helix play a role in stabilizing the trimer (Dev et al., 2016). In gp41CT_N, we have not observed any propensity for residues 707–710 to form an α helix, most likely because of the missing context of



Figure 3. Structural Features of the gp41CT_C Protein

(A) The C^{α} positional RMSDs between the 20 low-energy structures of gp41CT_C are represented as a sausage plot of the representative gp41CT_C structure (line thickness is proportional to the RMSD values). The RMSDs from an average structure were calculated after aligning each α -helical segment separately. For residues located at the overlapping boundaries of individual segments, an average of the two RMSD values was calculated. Bending between the helical fragments is variable in the 20 calculated structures.

(B) Cartoon representation of the $gp41CT_C$ protein showing the extensive hydrophobic interface formed by Leu, Ile, Val, Ala, Trp, and Phe residues (green sticks). (C) Surface representation of the $gp41CT_C$ colored according to electrostatic surface potential. A basic patch is formed in the arginine-rich LLP1 motif.

(D) Cartoon representation of the gp41CT_C protein showing Tyr and Trp residues as green sticks. Notice the cluster of six aromatic residues at the beginning of LLP3. The majority of aromatic residues are deeply buried in the interior of DPC micelle.

(E) Surface representation of the gp41CT_C protein showing Arg and Lys residues (blue spheres).

the transmembrane region. The C-terminal domain of gp41CT consists of three consecutive amphipathic α helices (LLP2, LLP3, and LLP1). By defining the boundaries of individual α helices, we are able to remove the long-standing uncertainty about the exact topology of the LLP segments. Our data revealed variable degree of membrane penetration among the three helices with the N-terminal LLP2 helix penetrating deeper than LLP3 and LLP1 (Figure 5). We have also shown that the helical structures of LLP2 and LLP3 contain several cation- π interactions between aromatic and basic residues in the *i* and i + 4 positions, respectively. Cation- π interacting pairs tend not to be buried while at the same time they are not as exposed as cationic residues alone (Gallivan and Dougherty, 2000). As pointed out previously (Steckbeck et al., 2011), HIV-1 gp41CT_C contains a number of highly conserved arginines and lysines within the LLP1 and LLP2 motifs. Our results show that several of those residues are involved in cation- π interactions. Analysis of over 5,000 Env sequences of HIV-1 M group isolates in the Los Alamos database revealed that those pairs are highly conserved (Figure S5). The unusually high number and conservation of these interactions suggest that they play a role in the stabilization of gp41CT_C a-helical structure. We speculate that they may also contribute to bilayer destabilization by keeping the charged side chain in the hydrophobic interior of PM or vice versa, exposing the bulky aromatic residues to the polar membrane interface. LLP peptides derived from the gp41CT protein were shown to disrupt membranes at very low concentrations (Costin et al., 2007). Interestingly, protein sequence alignment suggests that cation- π interactions are less prominent in HIV-2 and SIV strains (Figure S6). Several arginine residues are also highly conserved across the gp41CT sequence in 128 HIV-2 strains, but much less so in 206 SIV strains. However, an arginine-rich segment (IPRRIRQG₈₅₀) in LLP1 appears to be almost strictly conserved among all HIV-1, HIV-2, and SIV isolates (Figures S5 and S6). We found that this motif is structurally flexible, which suggests that it may have a functional role in gp41CT-mediated



processes. Interestingly, sequence analysis of EIAV gp41CT (GenBank: M16575.1), the longest among retroviruses (200 amino acids), revealed almost no sequence identity to HIV-1, HIV-2, or SIV, and appears to lack the LLP1 Arg-rich domain and frequent cation- π interactions. In summary, the structure of HIV-1 gp41CT revealed novel features that may be important for its function.

Early studies reported that antiserum produced against a synthetic peptide from gp41CT (residues 728–745) bound to HIV-1 Env, and that serum from HIV-1-infected humans also recognized this synthetic peptide (Kennedy et al., 1986). This observation led to the suggestion that this sequence is exposed on the virion surface to allow antibody binding and neutralization. Similarly, studies by Lu et al. (2008) supported the transient exposure of LLP2 on the cell surface during cell-cell fusion. In contrast, it was reported that antibodies that bind to the Kennedy sequence do not bind Env on intact virions (Steckbeck et al., 2010). In an attempt to explain these observations, Dimmock and colleagues proposed an alternative model by which the KE is exposed

Figure 4. Intermolecular Contacts Between gp41CT_C and Micelles

(A) 3D ¹³C-half-filtered/¹³C-edited NOESY spectrum of ¹³C-labeled gp41CT_C showing intermolecular NOEs between DPC and gp41CT_C residues. Colored intermolecular NOE crosspeaks correspond to colored atoms on the DPC structure.

(B) A selected region of 2D 1 H- 1 H NOESY spectrum of gp41CT_c showing NOEs between side chains of aromatic residues and DPC acyl chain methylene groups.

(Cleveland et al., 2003; Hollier and Dimmock, 2005). In this model, a three β -sheet membrane-spanning domain (MSD) was proposed based on theoretical sequence analysis. Our data revealed that the recombinant gp41CT_N protein does not contain an MSD or any secondary structure in the free and micelle-bound forms. Therefore, at least *in vitro*, our structural findings support a topology by which gp41CT_N is unstructured and not membrane associated (Figure 5).

Several models have been proposed to explain Env incorporation into virus particles (Checkley et al., 2011). The "direct Gag-gp41CT interaction" model is based on genetic and limited biochemical data (Cosson, 1996; Dorfman et al., 1994; Freed and Martin, 1995, 1996; Murakami and Freed, 2000a; Yu et al., 1992). It was shown that mutations in MA (L13E and L31E) block the incorporation of Env into HIV-1 particles (Freed and Martin, 1995). This phenotype, however, is reversed by truncating the gp41CT by 104 or 144

amino acids (Freed and Martin, 1995). Subsequent studies revealed that incorporation into virions of Env lacking the last 23, 30, 51, or 56 residues from the CT domain is specially blocked by MA L13E mutation, whereas truncations greater than 93 amino acids reverse this defect (Freed and Martin, 1996). Recently, a model has been proposed whereby MA hexamers of trimers form a lattice capable of accommodating gp41CT (Tedbury et al., 2016). According to our structural data, the region spanning LLP2-LLP3, which has been implicated in MA interaction (Murakami and Freed, 2000a), is the most membrane-sequestered region of gp41CT (Figure 5). Therefore, we propose a potential link between the degree of membrane association of gp41CT and its function in mediating Env incorporation. Because MA binding to the inner leaflet of the membrane is also required for Gag assembly, docking of the two proteins on a specific PM compartment is perhaps needed for a potential interaction to occur. This model, which is called "Gag-Env cotargeting" was among those proposed previously (Checkley et al., 2011). Based on previous studies and the



Figure 5. Membrane Interaction of $gp41CT_{c}$ and Overall Env Organization on the Virion Surface

(A) A model of gp41CT_C bound to a membrane bilayer constructed based on the NMR data using the representative structure of gp41CT_C with only minor modifications of the dihedral angles in the hinge regions to create an extended molecule. Length of the extended gp41CT_C domain shown here is 160 Å. Top and bottom panels show side and top views of the protein, respectively. Residues indicated as red spheres interact extensively with the interior of the membrane while those in blue are mostly exposed and interact with the polar head. The membrane bilayer was generated with VMD membrane builder plug-in (Humphrey et al., 1996).

(B) Top panel: A model depicting the gp120 and gp41 proteins on the surface of HIV-1 particles. The gp41CT_C domain is penetrating deeply in the inner leaflet of the membrane. Lower panel: An expanded view of the inner leaflet of the membrane showing gp41CT penetrating the bilayer.

current work, other scenarios for MA-gp41CT interaction are still possible: the gp41CT_N loop may play a role in MA interaction, and gp41CT can adopt multiple conformations depending on whether Gag is present and/or whether the underlying Gag lattice is in the immature or mature state. The advancement achieved here will allow for the development of new ap-

proaches to characterize the interplay between gp41CT protein and Gag.

In the "indirect interaction model," it was proposed that gp41CT-Gag interaction is mediated by cellular factor(s) (Checkley et al., 2011). This model is supported by the finding that the requirement for gp41CT in Env incorporation is celltype dependent (Akari et al., 2000; Murakami and Freed, 2000b). Numerous cellular factors have been described to interact with gp41CT, but their exact functional role is not well understood (reviewed in Santos da Silva et al., 2013). Nonetheless, in most cases, these interactions enhance virus replication. Among the cellular proteins most implicated in Env trafficking and incorporation are the adaptor protein complexes, AP-1 and AP-2 (Berlioz-Torrent et al., 1999; Boge et al., 1998; Ohno et al., 1997). These complexes are known to direct the sorting and trafficking of proteins in the secretory and endocytic pathway. AP-1 has been shown to regulate the subcellular localization of Env via binding to a dileucine motif (L855-L856) in gp41CT (Berlioz-Torrent et al., 1999; Byland et al., 2007; Wyss et al., 2011). AP-2, on the other hand, has been shown to drive clathrin-mediated endocytosis of Env from the PM by binding to gp41CT via a conserved Yxx (() indicates a hydrophobic residue; Y712SPL in our sequence) motif located at the N terminus of gp41CT (Boge et al., 1998; Ohno et al., 1997). Interestingly, AP-1 and AP-2 have also been shown to interact with the MA domain of Gag (Batonick et al., 2005; Camus et al., 2007). Our structural data show that the YSPL motif is unstructured and not membrane bound; the dileucine motif, which is located at the end of the gp41CT protein (Figure 3B), interacts only weakly with the membrane. Therefore, both motifs are likely accessible to interact with cellular factors. A recent structural study revealed that, even in the presence of the transmembrane segment, the YSPL motif is disordered (Dev et al., 2016). Whereas the YSPL motif and L855 are strictly conserved in HIV-1 isolates, L856 is more variable (Figure S5). In addition to those two motifs, it has been shown that Y795/W796 is critical in mediating cell-type-dependent Env incorporation by interacting with the FIP1C protein (Qi et al., 2013, 2015). Our structural data revealed that while W796 is buried and interacts with the membrane, Y795 is exposed to the cytoplasm and accessible for interactions with cellular factors (Figures 2B and 3D). It has yet to be determined how these motifs are involved in Env incorporation.

It has been suggested that part of MPER helices is inserted in the viral membrane, which would induce the membrane curvature required for fusion (Buzon et al., 2010). Our finding that the majority of gp41CT protein is embedded in the membrane, and the fact that gp41CT LLP peptides are able to disrupt membranes (Costin et al., 2007), support the hypothesis that gp41CT may contribute additional free energy to the fusion reaction. Indeed, it has been previously shown that during HIV-1 fusion, LLP2 may be transiently exposed on the surface of the effector cell in the presence of the target cell (Lu et al., 2008). A synthetic peptide derived from the LLP2 sequence was shown to interact with the gp41 core, suggesting that gp41CT may regulate the fusogenicity of viral Env through the interaction of the transiently exposed LLP2 region with the gp41 core on virions or HIV-1-infected cells (Lu et al., 2008).

This work may also help advance the current understanding of the structure, function, mobility, and lateral movement of Env on the virion surface. The mobility of Env spikes on the virion surface has been of high interest, as motion between these spikes might allow for Env clustering, receptor binding, and membrane fusion, and even IgG avidity (Sougrat et al., 2007). By employing electron tomographic methods, it has been shown that HIV-1 and SIV viruses make contact with T cells via a unique structure called viral "entry claw," which is typically composed of about six clustered rods of density that span the contact region (Sougrat et al., 2007). The finding that spacing of these rods is slightly closer than the average spacing of Env spikes observed in free viruses led to the suggestion that rearrangement of spikes within the viral membrane may be involved in formation of the structure. Interestingly, Env clustering has been recently recapitulated upon murine leukemia virus binding to the host cell (Riedel et al., 2017). Based on these observations, it is reasonable to hypothesize that the largely membrane-embedded HIV-1 gp41CT domain may facilitate or, at least does not constrain, clustering and lateral movement of Env spikes on the virion surface upon binding to host receptor.

Recent studies have shown that unliganded HIV-1 Env is intrinsically dynamic, transitioning between three distinct pre-fusion conformations, whose relative occupancies were remodeled by CD4 and antibody binding (Munro et al., 2014). CT truncation affected the antigenic surface of the ectodomain of HIV-1 Env on the opposite side of the PM (Chen et al., 2015). It was suggested that a physical coupling (conformation and/or dynamics) between the CT and the ectodomain is mediated by the transmembrane domain (Dev et al., 2016). In the absence of a detailed molecular model of the intact Env including gp41CT, it is difficult to understand this coupling relationship and how the CT domain can influence Env conformation and behavior to play a role in antibody recognition. We believe that our findings may help in the development of a platform to produce the intact Env protein, which may ultimately aid in developing new approaches for antiviral drug design, broadly neutralizing antibodies, and vaccine candidates.

STAR***METHODS**

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- CONTACT FOR REAGENT AND RESOURCE SHARING
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
- Bacterial Strains
- METHOD DETAILS
 - Plasmid Construction
 - Preparation of the gp41CT Protein
 - Protein Expression and Purification
 - Biochemical and Biophysical Characterization of the gp41CT Protein
 - Gel Filtration Assay
 - Circular Dichroism (CD) Spectroscopy
 - NMR Spectroscopy
 - O Paramagnetic Relaxation Enhancement NMR
 - Structure Calculations
- DATA AND SOFTWARE AVAILABILITY

SUPPLEMENTAL INFORMATION

Supplemental Information includes six figures and can be found with this article online at https://doi.org/10.1016/j.str.2017.09.010.

AUTHOR CONTRIBUTIONS

Conceptualization, R.E.M., A.B.S., J.V., and J.S.S.; Methodology, R.E.M., A.B.S., J.V., and J.S.S.; Investigation, R.E.M., A.B.S., and J.V.; Writing – Original Draft, R.E.M., J.V., and J.S.S.; Writing – Review & Editing, R.E.M., J.V., and J.S.S.; Funding Acquisition, J.S.S.; Resources, J.S.S.; Supervision, J.S.S.

ACKNOWLEDGMENTS

This work was supported by the National Institute of Health (R01 GM117837) and the Center for AIDS Research at the University of Alabama at Birmingham (P30 AI027767).

Received: July 8, 2017 Revised: August 17, 2017 Accepted: September 18, 2017 Published: October 19, 2017

REFERENCES

Akari, H., Fukumori, T., and Adachi, A. (2000). Cell-dependent requirement of human immunodeficiency virus type 1 gp41 cytoplasmic tail for Env incorporation into virions. J. Virol. *74*, 4891–4893.

Alam, S.M., Morelli, M., Dennison, S.M., Liao, H.X., Zhang, R., Xia, S.M., Rits-Volloch, S., Sun, L., Harrison, S.C., Haynes, B.F., et al. (2009). Role of HIV membrane in neutralization by two broadly neutralizing antibodies. Proc. Natl. Acad. Sci. USA *106*, 20234–20239.

Anraku, K., Fukuda, R., Takamune, N., Misumi, S., Okamoto, Y., Otsuka, M., and Fujita, M. (2010). Highly sensitive analysis of the interaction between HIV-1 Gag and phosphoinositide derivatives based on surface plasmon resonance. Biochemistry *49*, 5109–5116.

Baker, N.A., Sept, D., Joseph, S., Holst, M.J., and McCammon, J.A. (2001). Electrostatics of nanosystems: application to microtubules and the ribosome. Proc. Natl. Acad. Sci. USA *98*, 10037–10041.

Batonick, M., Favre, M., Boge, M., Spearman, P., Höning, S., and Thali, M. (2005). Interaction of HIV-1 Gag with the clathrin-associated adaptor AP-2. Virology *342*, 190–200.

Berlioz-Torrent, C., Shacklett, B.L., Erdtmann, L., Delamarre, L., Bouchaert, I., Sonigo, P., Dokhelar, M.C., and Benarous, R. (1999). Interactions of the cytoplasmic domains of human and simian retroviral transmembrane proteins with components of the clathrin adaptor complexes modulate intracellular and cell surface expression of envelope glycoproteins. J. Virol. 73, 1350–1361.

Boge, M., Wyss, S., Bonifacino, J.S., and Thali, M. (1998). A membrane-proximal tyrosine-based signal mediates internalization of the HIV-1 envelope glycoprotein via interaction with the AP-2 clathrin adaptor. J. Biol. Chem. *273*, 15773–15778.

Boscia, A.L., Akabori, K., Benamram, Z., Michel, J.A., Jablin, M.S., Steckbeck, J.D., Montelaro, R.C., Nagle, J.F., and Tristram-Nagle, S. (2013). Membrane structure correlates to function of LLP2 on the cytoplasmic tail of HIV-1 gp41 protein. Biophys. J. *105*, 657–666.

Buzon, V., Natrajan, G., Schibli, D., Campelo, F., Kozlov, M.M., and Weissenhorn, W. (2010). Crystal structure of HIV-1 gp41 including both fusion peptide and membrane proximal external regions. PLoS Pathog. *6*, e1000880.

Byland, R., Vance, P.J., Hoxie, J.A., and Marsh, M. (2007). A conserved dileucine motif mediates clathrin and AP-2-dependent endocytosis of the HIV-1 envelope protein. Mol. Biol. Cell *18*, 414–425.

Camus, G., Segura-Morales, C., Molle, D., Lopez-Vergès, S., Begon-Pescia, C., Cazevieille, C., Schu, P., Bertrand, E., Berlioz-Torrent, C., and Basyuk, E. (2007). The clathrin adaptor complex AP-1 binds HIV-1 and MLV Gag and facilitates their budding. Mol. Biol. Cell *18*, 3193–3203.

Checkley, M.A., Luttge, B.G., and Freed, E.O. (2011). HIV-1 envelope glycoprotein biosynthesis, trafficking, and incorporation. J. Mol. Biol. *410*, 582–608.

Chen, J., Kovacs, J.M., Peng, H., Rits-Volloch, S., Lu, J., Park, D., Zablowsky, E., Seaman, M.S., and Chen, B. (2015). HIV-1 ENVELOPE. Effect of the cytoplasmic domain on antigenic characteristics of HIV-1 envelope glycoprotein. Science 349, 191–195.

Chukkapalli, V., and Ono, A. (2011). Molecular determinants that regulate plasma membrane association of HIV-1 gag. J. Mol. Biol. *410*, 512–524.

Cleveland, S.M., McLain, L., Cheung, L., Jones, T.D., Hollier, M., and Dimmock, N.J. (2003). A region of the C-terminal tail of the gp41 envelope glycoprotein of human immunodeficiency virus type 1 contains a neutralizing epitope: evidence for its exposure on the surface of the virion. J. Gen. Virol. *84*, 591–602.

Cosson, P. (1996). Direct interaction between the envelope and matrix proteins of HIV-1. EMBO J. *15*, 5783–5788.

Costin, J.M., Rausch, J.M., Garry, R.F., and Wimley, W.C. (2007). Viroporin potential of the lentivirus lytic peptide (LLP) domains of the HIV-1 gp41 protein. Virol. J. 4, 123.

Delaglio, F., Grzesiek, S., Vuister, G.W., Zhu, G., Pfeifer, J., and Bax, A. (1995). NMRPipe: a multidimensional spectral processing system based on UNIX pipes. J. Biomol. NMR *6*, 277–293.

Dev, J., Park, D., Fu, Q., Chen, J., Ha, H.J., Ghantous, F., Herrmann, T., Chang, W., Liu, Z., Frey, G., et al. (2016). Structural basis for membrane anchoring of HIV-1 envelope spike. Science *353*, 172–175.

Dolinsky, T.J., Nielsen, J.E., McCammon, J.A., and Baker, N.A. (2004). PDB2PQR: an automated pipeline for the setup of Poisson-Boltzmann electrostatics calculations. Nucleic Acids Res. *32*, W665–W667.

Dorfman, T., Mammano, F., Haseltine, W.A., and Göttlinger, H.G. (1994). Role of the matrix protein in the virion association of the human immunodeficiency virus type 1 envelope glycoprotein. J. Virol. *68*, 1689–1696.

Freed, E.O. (2015). HIV-1 assembly, release and maturation. Nat. Rev. Microbiol. 13, 484–496.

Freed, E.O., and Martin, A.M. (1996). Domains of the human immunodeficiency virus type 1 matrix and gp41 cytoplasmic tail required for envelope incorporation into virions. J. Virol. 70, 341–351.

Freed, O.E., and Martin, A.M. (1995). Virion incorporation of envelope glycoproteins with long but not short cytoplasmic tails is blocked by specific, single amino acid substitutions in the human immunodeficiency virus type 1 matrix. J. Virol. 69, 1984–1989.

Gallivan, J.P., and Dougherty, D.A. (1999). Cation-pi interactions in structural biology. Proc. Natl. Acad. Sci. USA *96*, 9459–9464.

Gallivan, J.P., and Dougherty, D.A. (2000). A computational study of cation $-\pi$ interactions vs salt bridges in aqueous media: implications for protein engineering. J. Am. Chem. Soc. *122*, 870–874.

Güntert, P. (2004). Automated NMR structure calculation with CYANA. Methods Mol. Biol. *278*, 353–378.

Hales, C.M., Griner, R., Hobdy-Henderson, K.C., Dorn, M.C., Hardy, D., Kumar, R., Navarre, J., Chan, E.K., Lapierre, L.A., and Goldenring, J.R. (2001). Identification and characterization of a family of Rab11-interacting proteins. J. Biol. Chem. *276*, 39067–39075.

Hollier, M.J., and Dimmock, N.J. (2005). The C-terminal tail of the gp41 transmembrane envelope glycoprotein of HIV-1 clades A, B, C, and D may exist in two conformations: an analysis of sequence, structure, and function. Virology 337, 284–296.

Humphrey, W., Dalke, A., and Schulten, K. (1996). VMD: visual molecular dynamics. J. Mol. Graph. 14, 33–38, 27–28.

Jarvet, J., Zdunek, J., Damberg, P., and Graslund, A. (1997). Three-dimensional structure and position of porcine motilin in sodium dodecyl sulfate micelles determined by 1H NMR. Biochemistry *36*, 8153–8163.

Johnson, B.A., and Blevins, R.A. (1994). NMRview: a computer program for the visualization and analysis of NMR data. J. Biomol. NMR *4*, 603–614.

Kennedy, R.C., Henkel, R.D., Pauletti, D., Allan, J.S., Lee, T.H., Essex, M., and Dreesman, G.R. (1986). Antiserum to a synthetic peptide recognizes the HTLV-III envelope glycoprotein. Science *231*, 1556–1559.

Kim, Y.P., Yeo, K.J., Kim, M.H., Kim, Y.C., and Jeon, Y.H. (2010). Structural characterization of the intra-membrane histidine kinase YbdK from Bacillus subtilis in DPC micelles. Biochem. Biophys. Res. Commun. *391*, 1506–1511.

Kuhlmann, A.S., Steckbeck, J.D., Sturgeon, T.J., Craigo, J.K., and Montelaro, R.C. (2014). Unique functional properties of conserved arginine residues in the lentivirus lytic peptide domains of the C-terminal tail of HIV-1 gp41. J. Biol. Chem. *289*, 7630–7640.

Lambele, M., Labrosse, B., Roch, E., Moreau, A., Verrier, B., Barin, F., Roingeard, P., Mammano, F., and Brand, D. (2007). Impact of natural polymorphism within the gp41 cytoplasmic tail of human immunodeficiency virus type 1 on the intracellular distribution of envelope glycoproteins and viral assembly. J. Virol. *81*, 125–140.

Laskowski, R.A., Rullmann, J.A.C., MacArthur, M.W., Kaptein, R., and Thornton, J.M. (1996). AQUA and Procheck NMR: programs for checking the quality of protein structures solved by NMR. J. Biomol. NMR 8, 477–486.

Lazaridis, T., Mallik, B., and Chen, Y. (2005). Implicit solvent simulations of DPC micelle formation. J. Phys. Chem. B *109*, 15098–15106.

Lu, L., Zhu, Y., Huang, J., Chen, X., Yang, H., Jiang, S., and Chen, Y.H. (2008). Surface exposure of the HIV-1 env cytoplasmic tail LLP2 domain during the membrane fusion process: interaction with gp41 fusion core. J. Biol. Chem. 283, 16723–16731.

Markley, J.L., Bax, A., Arata, Y., Hilbers, C.W., Kaptein, R., Sykes, B.D., Wright, P.E., and Wuthrich, K. (1998). Recommendations for the presentation of NMR structures of proteins and nucleic acids. J. Mol. Biol. 280, 933–952.

Mercredi, P.Y., Bucca, N., Loeliger, B., Gaines, C.R., Mehta, M., Bhargava, P., Tedbury, P.R., Charlier, L., Floquet, N., Muriaux, D., et al. (2016). Structural and molecular determinants of membrane binding by the HIV-1 matrix protein. J. Mol. Biol. *428*, 1637–1655.

Merk, A., and Subramaniam, S. (2013). HIV-1 envelope glycoprotein structure. Curr. Opin. Struct. Biol. 23, 268–276.

Munro, J.B., Gorman, J., Ma, X., Zhou, Z., Arthos, J., Burton, D.R., Koff, W.C., Courter, J.R., Smith, A.B., 3rd, Kwong, P.D., et al. (2014). Conformational dynamics of single HIV-1 envelope trimers on the surface of native virions. Science *346*, 759–763.

Murakami, T., and Freed, E.O. (2000a). Genetic evidence for an interaction between human immunodeficiency virus type 1 matrix and α -helix 2 of the gp41 cytoplasmic tail. J. Virol. 74, 3548–3554.

Murakami, T., and Freed, E.O. (2000b). The long cytoplasmic tail of gp41 is required in a cell type-dependent manner for HIV-1 envelope glycoprotein incorporation into virions. Proc. Natl. Acad. Sci. USA 97, 343–348.

Muranyi, W., Malkusch, S., Müller, B., Heilemann, M., and Kräusslich, H.G. (2013). Super-resolution microscopy reveals specific recruitment of HIV-1 envelope proteins to viral assembly sites dependent on the envelope C-terminal tail. PLoS Pathog. *9*, e1103198.

Ohno, H., Aguilar, R.C., Fournier, M.C., Hennecke, S., Cosson, P., and Bonifacino, J.S. (1997). Interaction of endocytic signals from the HIV-1 envelope glycoprotein complex with members of the adaptor medium chain family. Virology *238*, 305–315.

Ono, A., Ablan, S.D., Lockett, S.J., Nagashima, K., and Freed, E.O. (2004). Phosphatidylinositol (4,5) bisphosphate regulates HIV-1 Gag targeting to the plasma membrane. Proc. Natl. Acad. Sci. USA *101*, 14889–14894.

Papavoine, C.H., Konings, R.N., Hilbers, C.W., and van de Ven, F.J. (1994). Location of M13 coat protein in sodium dodecyl sulfate micelles as determined by NMR. Biochemistry *33*, 12990–12997.

Qi, M., Chu, H., Chen, X., Choi, J., Wen, X., Hammonds, J., Ding, L., Hunter, E., and Spearman, P. (2015). A tyrosine-based motif in the HIV-1 envelope glycoprotein tail mediates cell-type- and Rab11-FIP1C-dependent incorporation into virions. Proc. Natl. Acad. Sci. USA *112*, 7575–7580.

Qi, M., Williams, J.A., Chu, H., Chen, X., Wang, J.J., Ding, L., Akhirome, E., Wen, X., Lapierre, L.A., Goldenring, J.R., et al. (2013). Rab11-FIP1C and

Rab14 direct plasma membrane sorting and particle incorporation of the HIV-1 envelope glycoprotein complex. PLoS Pathog. 9, e1003278.

Riedel, C., Vasishtan, D., Siebert, C.A., Whittle, C., Lehmann, M.J., Mothes, W., and Grunewald, K. (2017). Native structure of a retroviral envelope protein and its conformational change upon interaction with the target cell. J. Struct. Biol. *197*, 172–180.

Roy, N.H., Chan, J., Lambele, M., and Thali, M. (2013). Clustering and mobility of HIV-1 Env at viral assembly sites predict its propensity to induce cell-cell fusion. J. Virol. *87*, 7516–7525.

Saad, J.S., Miller, J., Tai, J., Kim, A., Ghanam, R.H., and Summers, M.F. (2006). Structural basis for targeting HIV-1 Gag to virus assembly sites on the plasma membrane. Proc. Natl. Acad. Sci. USA *103*, 11364–11369.

Samal, A.B., Ghanam, R.H., Fernandez, T.F., Monroe, E.B., and Saad, J.S. (2011). NMR, biophysical and biochemical studies reveal the minimal calmodulin-binding domain of the HIV-1 matrix protein. J. Biol. Chem. 286, 33533–33543.

Santos da Silva, E., Mulinge, M., and Perez Bercoff, D. (2013). The frantic play of the concealed HIV envelope cytoplasmic tail. Retrovirology *10*, 54.

Sham, S.W., McDonald, J.M., Micoli, K.J., and Krishna, N.R. (2008). Solution structure of a calmodulin-binding domain in the carboxy-terminal region of HIV type 1 gp160. AIDS Res. Hum. Retroviruses *24*, 607–616.

Shen, Y., Delaglio, F., Cornilescu, G., and Bax, A. (2009). TALOS+: a hybrid method for predicting protein backbone torsion angles from NMR chemical shifts. J. Biomol. NMR *44*, 213–223.

Shkriabai, N., Datta, S.A., Zhao, Z., Hess, S., Rein, A., and Kvaratskhelia, M. (2006). Interactions of HIV-1 Gag with assembly cofactors. Biochemistry *45*, 4077–4083.

Sougrat, R., Bartesaghi, A., Lifson, J.D., Bennett, A.E., Bess, J.W., Zabransky, D.J., and Subramaniam, S. (2007). Electron tomography of the contact between T cells and SIV/HIV-1: implications for viral entry. PLoS Pathog. *3*, e63.

Srinivas, S.K., Srinivas, R.V., Anantharamaiah, G.M., Segrest, J.P., and Compans, R.W. (1992). Membrane interactions of synthetic peptides corresponding to amphipathic helical segments of the human immunodeficiency virus type-1 envelope glycoprotein. J. Biol. Chem. *267*, 7121–7127.

Steckbeck, J.D., Craigo, J.K., Barnes, C.O., and Montelaro, R.C. (2011). Highly conserved structural properties of the C-terminal tail of HIV-1 gp41 protein

despite substantial sequence variation among diverse clades: implications for functions in viral replication. J. Biol. Chem. 286, 27156–27166.

Steckbeck, J.D., Kuhlmann, A.S., and Montelaro, R.C. (2013). C-terminal tail of human immunodeficiency virus gp41: functionally rich and structurally enigmatic. J. Gen. Virol. *94*, 1–19.

Steckbeck, J.D., Sun, C., Sturgeon, T.J., and Montelaro, R.C. (2010). Topology of the C-terminal tail of HIV-1 gp41: differential exposure of the Kennedy epitope on cell and viral membranes. PLoS One 5, e15261.

Tedbury, P.R., Novikova, M., Ablan, S.D., and Freed, E.O. (2016). Biochemical evidence of a role for matrix trimerization in HIV-1 envelope glycoprotein incorporation. Proc. Natl. Acad. Sci. USA *113*, E182–E190.

Ulmer, T.S., and Bax, A. (2005). Comparison of structure and dynamics of micelle-bound human alpha-synuclein and Parkinson disease variants. J. Biol. Chem. *280*, 43179–43187.

Ulmer, T.S., Bax, A., Cole, N.B., and Nussbaum, R.L. (2005). Structure and dynamics of micelle-bound human alpha-synuclein. J. Biol. Chem. *280*, 9595–9603.

Vlach, J., and Saad, J.S. (2013). Trio engagement via plasma membrane phospholipids and the myristoyl moiety governs HIV-1 matrix binding to bilayers. Proc. Natl. Acad. Sci. USA *110*, 3525–3530.

Vranken, W.F., Boucher, W., Stevens, T.J., Fogh, R.H., Pajon, A., Llinas, M., Ulrich, E.L., Markley, J.L., Ionides, J., and Laue, E.D. (2005). The CCPN data model for NMR spectroscopy: development of a software pipeline. Proteins 59, 687–696.

Wyss, S., Berlioz-Torrent, C., Boge, M., Blot, G., Höning, S., Benarous, R., and Thali, M. (2011). The highly conserved C-terminal dileucine motif in the cytosolic domain of the human immunodeficiency virus type 1 envelope glycoprotein is critical for its association with the AP-1 clathrin adaptor [correction of adapter]. J. Virol. *75*, 2982–2992.

Yu, X., Yuan, X., Matsuda, Z., Lee, T.-H., and Essex, M. (1992). The matrix protein of human immunodeficiency virus type I is required for incorporation of viral envelope protein into mature virions. J. Virol. *66*, 4966–4971.

Yuan, T., Mietzner, T.A., Montelaro, R.C., and Vogel, H.J. (1995). Characterization of the calmodulin binding domain of SIV transmembrane glycoprotein by NMR and CD spectroscopy. Biochemistry *34*, 10690–10696.

STAR***METHODS**

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and Virus Strains		
BL21-CodonPlus (DE3)-RIL Competent Cells	Agilent Technology	Cat#230245
XL 10-Gold Ultracompetent Cells	Agilent Technology	Cat#200517-4
Subcloning Efficiency DH5a Competent Cells	ThermoFisher	Cat#18265017
HIV-1 recombinant clone pNL4-3	NIH AIDS reagents	Cat#114
Chemicals, Peptides, and Recombinant Proteins		
SUMO-protease	(Samal et al., 2011)	N/A
Lysozyme	Sigma-Aldrich	Cat#L2879
Enzyme Xhol	BioLabs	Cat#R0146S
Enzyme BamHI-HF	BioLabs	Cat#R3136S
n-dodecylphosphocholine (DPC)	Avanti Polar Lipids	Cat#850336P
n-dodecylphosphocholine (DPC)	Bio and Beyond	Fos-Choline-12 (DPC)
1,2-dihexanoyl-sn-glycero-3-phosphocholine (DHPC)	Avanti Polar Lipids	Cat#850305P
1,2-dimyristoyl-sn-glycero-3-phosphocholine (DMPC)	Avanti Polar Lipids	Cat#860345P
16-DOXYL-stearic acid (16-DSA)	Sigma-Aldrich	Cat#253596
5-DOXYL-stearic acid (5-DSA)	Sigma-Aldrich	Cat#253634
Ammonium- ¹⁵ N chloride (¹⁵ NH ₄ Cl)	Sigma-Aldrich	Cat#299251
D-glucose- ¹³ C ₆	Sigma-Aldrich	Cat#389374
Deuterium oxide (D ₂ O)	Sigma-Aldrich	Cat#151882
Gadodiamide	Selleckchem	Cat#S1751
Minimum essential medium (MEM) vitamin solution	Gibco	Cat#11120052
Gel Filtration Calibration Kit LMW	GE Healthcare	Cat#28-4038-41
Phenylmethanesulfonyl fluoride	BACHEM	Cat#Q-1395
Benzamidine	ACROS Organics	Cat#105240050
Tris (2-Carboxyethyl) phosphine Hydrochloride (TCEP-HCl)	Gold Biotechnology	Cat#TCEP25
Isopropyl β-D-1-thiogalactopyranoside (IPTG)	Gold Biotechnology	Cat#I2481C
Triton X-100	ACROS Organics	Cat#215682500
Potassium fluoride	ACROS Organics	Cat#20135-2500
CutSmart Buffer	BioLabs	Cat#B7204S
Detergent screen kit	Hampton Research	Cat#HR2-408
Critical Commercial Assays		
QuikChange Lighting Site-Directed Mutagenesis Kit	Agilent Technologies	Cat#210518-5
illustra PuReTaq Ready-To-Go PCR Beads	GE Healthcare Life Science	Cat#27-9559-01
Deposited Data		
Coordinates of gp41CT _C	PDB database	[PDB]: [5VWL]
NMR chemical shifts of gp41CT _N	BMRB database	[BMRB]: [27110]
NMR chemical shifts of gp41CT _C	BMRB database	[BMRB]: [30297]
Oligonucleotides		
Primers for full-length His ₆ -SUMO-gp41CT	Integrated DNA Technologies	https://www.idtdna.com/site
Primers for His ₆ -SUMO-gp41CT _N	Integrated DNA Technologies	https://www.idtdna.com/site
Primers for His ₆ -SUMO-gp41CT _C	Integrated DNA Technologies	https://www.idtdna.com/site
Recombinant DNA		
pET28a His ₆ -SUMO-gp41CT	This paper	N/A
pET28a His ₆ -SUMO-gp41CT _N	This paper	N/A

(Continued on next page)

Continued		
REAGENT or RESOURCE	SOURCE	IDENTIFIER
pET28a His ₆ -SUMO-gp41CT _C	This paper	N/A
Software and Algorithms		
PYMOL	Schrödinger	http://www.pymol.org/
NMRViewJ	(Johnson and Blevins, 1994)	http://www.onemoonscientific.com/ nmrviewj
CCPN Analysis	(Vranken et al., 2005)	http://www.ccpn.ac.uk
NMRPipe	(Delaglio et al., 1995)	https://spin.niddk.nih.gov/bax/ software/NMRPipe
TALOS+	(Shen et al., 2009)	https://spin.niddk.nih.gov/bax/ software/TALOS
CYANA	(Güntert, 2004)	http://www.cyana.org/wiki/ index.php/Main_Page
UGENE	Unipro	http://ugene.net
PROCHECK-NMR	(Laskowski et al., 1996)	http://www.ebi.ac.uk/thornton- srv/software/PROCHECK/
PDB2PQR	(Baker et al., 2001)	http://nbcr-222.ucsd.edu/ pdb2pqr_2.1.1/
APBS	(Dolinsky et al., 2004)	http://www.poissonboltzmann.org
wheel.pl	D. Armstrong and R. Zidovetzki (2001)	http://rzlab.ucr.edu/scripts/ wheel/wheel.html
Topspin	Bruker	https://www.bruker.com/
Image Lab	Bio-Rad	http://www.bio-rad.com/

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Jamil S. Saad (saad@uab.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Bacterial Strains

His₆-SUMO-gp41CT, His₆-SUMO-gp41CT_N, and His₆-SUMO-gp41CT_C proteins were overexpressed in *Escherichia coli* BL21 (DE3) codon plus RIL cells with Luria-Bertani (LB) broth or M9 minimal media supplemented with ¹⁵NH₄Cl and D-glucose-¹³C₆ and containing 50 μ g/mL Kanamycin.

METHOD DETAILS

Plasmid Construction

To construct the gp41CT clones, their coding sequences were PCR-amplified from the pNL4-3 isolate (NCBI accession code: AF324493) and ligated to the 3'-end of small ubiquitin-like modifier (SUMO) gene via BamHI and XhoI sites within a pET28 vector. The resulting construct has a His₆ tag encoded on the N-terminus of SUMO. Plasmids were verified by DNA sequencing at the Heflin Center for Genomic Sciences at the University of Alabama at Birmingham.

Preparation of the gp41CT Protein

The gp41CT protein was prone to proteolysis even during expression and purification. The proteolytic cleavage resulted in two fragments as detected by chromatography and SDS PAGE. Purification of the cleaved products yielded soluble (\sim 6 kDa) and insoluble (\sim 12 kDa) fragments. The small soluble fragment was isolated and identified by mass spectrometry as the N-terminal 45 amino acids (residues 707-751; gp41CT_N). The remaining insoluble C-terminal 105-amino acid region (residues 752-856) is referred to as gp41CT_C. We constructed two plasmids to express and purify the gp41CT_N and gp41CT_C proteins independently. Cleavage of SUMO tag leaves a non-native N-terminal serine residue on gp41CT and gp41CT_N proteins.

Protein Expression and Purification

gp41CT and gp41CT_c Proteins

To make uniformly ¹⁵N- and ¹⁵N-, ¹³C-labelled proteins, transformed cells were grown overnight in 25 mL LB medium (50 mg/L kanamycin) at 37°C. Cells were then spun down and transferred into 1 L of modified M9 minimal medium containing ¹⁵NH₄Cl and D-glucose-¹³C₆ as the sole nitrogen and carbon sources to produce ¹⁵N- and ¹³C-labelled proteins, respectively. The M9 minimal medium contained 6.8 g/L Na₂HPO₄, 3 g/L KH₂PO₄ (pH 7.4), 0.5 g/L NaCl, 5 mM MgSO₄, 0.2 mM CaCl₂, 0.25 mL/L trace metal solution (1000x), 2.5 mL/L minimum essential medium (MEM) vitamin solution, 1 g/L ¹⁵NH₄Cl, 2 g/L ¹³C D-glucose, and 50 mg/L kanamycin. Cells were grown at 37°C until A_{600} reached 1.6-1.8, then induced with 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) and grown at 37°C for 2 h. Cells were then spun down and stored overnight at -80°C. Next day, the cell pellet was resuspended in lysis buffer containing 100 mM sodium phosphates (pH 8.0), 500 mM NaCl, 0.5% Triton X-100, 10% glycerol, 20 mM imidazole, 0.5 mg/mL lysozyme, 2 mM benzamidine and 0.1 mM phenylmethanesulfonyl fluoride. Cells were sonicated and cell lysate was spun down at 35,000 g for 40 min at 4°C. The protein-containing supernatant was applied to cobalt affinity resin. The resin was washed with 10 column volumes of buffer A (20 mM sodium phosphates (pH 8.0), 500 mM NaCl, 10% glycerol, and 0.1% Triton X-100), buffer B (20 mM sodium phosphates (pH 8.0), 500 mM NaCl and 10% glycerol), and buffer C (20 mM sodium phosphates (pH 8.0), 500 mM NaCl and 10% glycerol), and buffer C (20 mM sodium phosphates (pH 8.0), 500 mM NaCl and 10% glycerol), and buffer C (20 mM sodium phosphates (pH 8.0), 500 mM NaCl and 10% glycerol. His₆-SUMO protease was then added to the resin and left to rock overnight at room temperature.

A detergent screen kit containing 96 unique conditions was used to elute gp41CT and gp41CT_C proteins from the cobalt resin, yielding twenty-three detergent conditions in which the proteins were partially or completely soluble. Among those detergents were dodecylphosphocholine (DPC), dihexanoylphosphatidylcholine (DHPC) and sodium dodecyl sulfate (SDS, Sigma-Aldrich) micelles. In addition, we were able to solubilize the protein in dimyristoylphosphatidylcholine (DMPC)/DHPC bicelles (prepared as described below). DPC was chosen for further work. The protein was eluted off the resin using 20 mM sodium phosphates (pH 8.0), 500 mM NaCl, 10% glycerol and 5.7 mM DPC. The fractions considered pure as judged by Coomassie-stained SDS-polyacrylamide gel electrophoresis (SDS-PAGE) were pooled, concentrated, and buffer was exchanged to 50 mM sodium phosphates (pH 6.0), 50 mM NaCl, 1 mM TCEP, and 5% D_2O (NMR buffer) by using Amicon Ultra Centrifugal Filters (3 kDa membrane cut-off). Final concentration of DPC in NMR samples was ~25 mM.

gp41CT_N Protein

To make uniformly ¹⁵N- and ¹⁵N-,¹³C-labelled samples, transformed cells were grown in 25 mL LB medium (50 mg/L kanamycin) at 37°C overnight. Cells were then spun down and transferred into 1 L M9 minimal medium containing ¹⁵NH₄Cl and D-glucose-¹³C₆ as the sole nitrogen and carbon sources to produce ¹⁵N- and ¹³C-labelled proteins, respectively. Cells were grown at 37°C until A_{600} was 0.8, induced with 1 mM IPTG and grown at 37°C overnight. Cells were spun down and stored overnight at -80°C. Next day, the cell pellet was resuspended in lysis buffer containing 100 mM sodium phosphates (pH 8.0), 500 mM NaCl, 2 mM benzamidine and 0.1 mM phenylmethanesulfonyl fluoride. Cells were sonicated and cell lysate was spun down at 35,000 g for 40 min at 4°C. The protein-containing supernatant was applied to cobalt affinity resin that was subsequently washed with 5 column volumes of a buffer containing 20 mM sodium phosphates (pH 8.0) and 300 mM NaCl. The His₆-SUMO-gp41CT_N protein was eluted using an imidazole gradient (0–300 mM). Fractions containing the protein were pooled and His₆-SUMO protease was added. Protein was dialyzed overnight at 4°C against a buffer containing 20 mM sodium phosphates (pH 8.0) and 300 mM NaCl. His₆-SUMO and His₆-SUMO protease were removed by cobalt affinity resin. The gp41CT_N fractions were pooled, concentrated and further purified by gel filtration chromatography. Fractions containing the gp41CT_N protein were then pooled and buffer was exchanged to NMR buffer by using Amicon Ultra Centrifugal Filters (3 kDa membrane cut-off).

Preparation of DMPC/DHPC Bicelles

Bicelles consisting of DMPC/DHPC in molar ratio (q) of 0.5 were prepared at 2 % (w/v) total lipid concentration by mixing appropriate amounts of solid DMPC with 336 mM DHPC in 20 mM sodium phosphates (pH 8.0), 500 mM NaCl and 10% glycerol buffer, followed by three freeze-thaw cycles. 2 mL of the bicelle solution was added to 1 mL cobalt resin containing His_6 -SUMO-gp41CT (or His_6 -SUMO-gp41CT_C) after imidazole removal (see above). His_6 -SUMO protease was then added and the resin was rocked overnight at room temperature. After separating the resin by filtration, gp41CT (or gp41CT_C) in bicelles was concentrated and buffer exchanged to NMR buffer using Amicon Ultra centrifugal filters. DHPC was added to the final sample to compensate for any loss during concentration. Final DHPC/DMPC ratio was verified by ¹H NMR.

Biochemical and Biophysical Characterization of the gp41CT Protein

NMR and CD spectroscopic methods were employed to identify the proper buffer conditions that maintained protein solubility, secondary structure, and most importantly, allowed for detection of NMR signals. Our initial attempts focused on properly reconstituting the protein in DMPC/DHPC bicelles. Unfortunately, the overall quality of NMR data was poor. Only a very small subset of ¹H-¹⁵N resonances were observed in the HSQC spectrum. The best results in terms of high solubility and protein stability were achieved with DPC micelles.

The oligomerization properties of gp41CT in DPC micelles were analyzed by a gel filtration assay and verified by SDS-PAGE (Figure S1). Protein elutes as a single peak at 14.8 mL (Figure S1), indicating a homogenous DPC-reconstituted protein. Interestingly, compared to known protein standards the apparent molecular mass of gp41CT protein in DPC micelle corresponds to a \sim 75 kDa species (Figure S1). Previous studies have shown that the size of a DPC micelle can range from 50 to 90 molecules which

corresponds to 18-32 kDa (Kim et al., 2010; Lazaridis et al., 2005). Therefore, the larger than expected estimated molecular weight is either due to the elongated shape of gp41CT and formation of a larger micelle to accommodate the protein or due to formation of a trimer in a smaller micelle. The latter has been ruled out as described below. Next, we examined the secondary structure of DPCassociated gp41CT protein by CD spectroscopy. The far-UV CD spectrum of gp41CT in DPC micelles shows two minima at 208 and 222 nm, features distinctive of an α -helical secondary structure (Figure S2). Our data are in agreement with previous CD studies conducted on short peptides derived from gp41CT that indicated the presence of α -helical structures in membranes consisting of 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC) and 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphoglycerol (POPG) (Costin et al., 2007) or SDS micelles (Steckbeck et al., 2011). For DPC-solubilized gp41CT, NMR spectra were collected at different temperatures with best data obtained at 50°C. However, numerous signals were relatively weak and severely overlapping which rendered signal assignments impractical (Figure S2). The CD spectra obtained for gp41CT in DPC micelles at 20 and 50°C (Figure S2) are virtually identical, demonstrating that the protein structure was not compromised at 50°C. In summary, none of the conditions applied to the gp41CT protein produced sufficient quality of NMR data to enable full structural characterization. This result is likely attributed to the presence of two independent domains (free and membrane-bound) with very different tumbling and relaxation rates.

Gel Filtration Assay

The mobility of $gp41CT_C$ was analyzed by a gel filtration assay. Briefly, 0.5 mL of ~200 μ M sample was loaded onto a HiLoad Superdex 75 (10/300) column in a buffer containing 50 mM sodium phosphates (pH 6), 50 mM NaCl, 1 mM TCEP, and 3 mM DPC. Protein fractions were analyzed by SDS-PAGE and stained by Coomassie brilliant blue. The approximate molecular weight of $gp41CT_C$ was determined by a molecular weight gel filtration calibration kit.

Circular Dichroism (CD) Spectroscopy

CD spectra were acquired on a Jasco J815 spectropolarimeter at 20, 35 and 50°C from 260 to 185 nm. Scanning rate was set to 50 nm/min. Loading concentrations were 20 μ M for gp41CT_N in a buffer containing 10 mM potassium phosphates (pH 6) and 100 mM potassium fluoride. The gp41CT_C was in a buffer containing 10 mM potassium phosphates (pH 6), 100 mM potassium fluoride and 3 mM DPC. The background signal from the buffer solution was subtracted from each spectrum. Protein samples were run on gel filtration column (as described above) to ensure high purity and homogeneity prior to collection of the CD spectra.

NMR Spectroscopy

NMR data were collected at 32 or 50°C on a Bruker Avance II (700 MHz¹H) spectrometer equipped with a cryogenic triple-resonance probe, processed with NMRPipe (Delaglio et al., 1995) and analyzed with NMRVIEW (Johnson and Blevins, 1994) or CCPN Analysis (Vranken et al., 2005). Isotopically unlabeled and uniformly ¹³C-, ¹⁵N-, or ¹³C-/¹⁵N-labelled gp41CT_N and gp41CT_C protein samples were prepared at ~400-500 μ M concentrations. The backbone and side-chain atom resonances were assigned using HNCA, HN(CO) CA, HNCACB, HN(CO)CACB, HNCO, HN(CA)CO, ¹⁵N-edited NOESY- and TOCSY-HSQC, and (H)CCH-TOCSY experiments. Assignments of aromatic signals for gp41CT_C protein were confirmed by the ¹H-¹³C TROSY, TROSY-(H)CCH-COSY and (HB) CB(CGCD)HD experiments. Intermolecular NOE contacts between DPC and gp41CT_C were observed in 2D NOESY, 3D ¹H-¹⁵N NOESY-HSQC, and ¹³C-half-filtered/¹³C-edited NOESY-HSQC spectra (all mixing times 120 ms). Chemical shifts were referenced to 2,2-dimethyl-2-silapentanesulfonic acid (Markley et al., 1998).

Paramagnetic Relaxation Enhancement NMR

Doxyl Stearic Acid Titrations

Solid aliquots of 5- or 16-doxyl stearic acid (5-DSA and 16-DSA, respectively) were added to 500 μ L of 150 μ M ¹⁵N-labelled gp41CT_C in NMR buffer and 25 mM DPC to final concentrations of 0.1, 0.2, 0.4, 0.8, and 1.6 mM. ¹H-¹⁵N HSQC spectra were collected at 50°C after each addition. The spin labels did not alter the protein structure given the absence of chemical shift perturbations.

Gadodiamide Titration

A stock solution of 100 mM gadodiamide in H₂O was titrated to 90 μ M ¹⁵N-labelled gp41CT_C in NMR buffer to final concentrations of 0.05, 0.1, 0.2, 0.4, 0.8, 1.6, 3.2, 6.4 and 9.6 mM. ¹H-¹⁵N HSQC spectra were collected at 50°C after each addition.

Data Analysis

For each titration, peak heights were determined for non-overlapping signals and relative reductions (RR) of the heights calculated according to RR = $(I_0 - I_n)/I_0$, where I_0 is peak height in the control experiment and I_n is peak height at titration point *n*.

Structure Calculations

Phi and psi dihedral angle restraints were generated in TALOS+ (Shen et al., 2009) based on ¹H^N, ¹⁵N, ¹³C', ¹H^{α}, ¹³C^{α} and ¹³C^{β} chemical shifts and used with an uncertainty of ±2 × standard deviation or ± 20 deg, whichever was larger, only for residues with a "Good" score (Shen et al., 2009). Structure of the gp41CT_C was calculated in CYANA (Güntert, 2004) using ϕ/ψ dihedral angles as predicted with TALOS+ and assigned NOE restraints from ¹H-¹⁵N NOESY-HSQC and ¹H-¹³C HMQC-NOESY. Upper distance limits were set to 2.7, 3.3, 4.0, 4.5 or 5.0 Å based on the intensities of NOE cross-peaks and standard pseudo atom distance corrections were applied to groups of degenerate hydrogen atoms during calculations. Hydrogen bond restraints with O–H and O–N distances of 1.8–2.0 and 2.7–3.0 Å, respectively, were used in α -helical regions identified by characteristic α -helical NOEs and TALOS prediction. Side chain rotamers were treated with low-weight torsion angle restraints during the initial stages of simulated annealing. Structure calculations

were initiated with random torsion angle values and standard protocols for simulated annealing and subsequent molecular dynamics minimization were used. In the final calculation cycle, 100 random input structures were minimized and 20 structures with the best agreement with the final restraint set were selected as a representative ensemble. Statistical information is provided in Table 1. The quality of structures was assessed using PROCHECK-NMR (Laskowski et al., 1996), and visualization was performed via PyMOL (The PyMOL Molecular Graphics System, Version 1.8.4 Schrödinger, LLC.). Electrostatic potential maps were generated using PDB2PQR and APBS software (Baker et al., 2001; Dolinsky et al., 2004).

DATA AND SOFTWARE AVAILABILITY

The atomic coordinates and restraints have been deposited in the Protein Data Bank, (PDB ID code 5VWL), and chemical shifts have been deposited in the Biological Magnetic Resonance Data Bank (accession codes 27110 and 30297 for $gp41CT_N$ and $gp41CT_C$, respectively).